ENTOMOLOGICAL NEWS

JULY 1961

Vol. LXXII **No. 7**

CONTENTS

Gillaspy—A new Stictiella from Mexico 169

McDermott—A new genus of firefly 174

Crabill—Concerning Neogeophilidae (cont.) 177

Selander—Meloid beetles from the West Indies 190

Judd—Melanagromyza from linden galls 192

Throne—Psectra diptera in Wisconsin 193

Review—A manual of common beetles 194

Rapp—Corrodentia in cliff swallow nests 195

PUBLISHED MONTHLY, EXCEPT AUGUST AND SEPTEMBER, BY
THE AMERICAN ENTOMOLOGICAL SOCIETY
PRINCE AND LEMON STS., LANCASTER, PA.
AND
1900 RACE STREET, PHILADELPHIA 3, PA.

Subscription, per yearly volume of ten numbers: $5.00 domestic; $5.30 foreign; $5.15 Canada.
Second-class postage paid at Lancaster, Pa.

ENTOMOLOGICAL NEWS

ENTOMOLOGICAL NEWS is published monthly, excepting August and September, by The American Entomological Society at Prince and Lemon Sts., Lancaster, Pa., and the Academy of Natural Sciences, 1900 Race Street, Philadelphia 3, Pa., U. S. A.

PHILIP P. CALVERT, Editor Emeritus. R. G. SCHMIEDER, Editor. Editorial Staff: H. J. GRANT, JR., E. J. F. MARX, M. E. PHILLIPS, and J. A. G. REHN.

SUBSCRIPTIONS: Communications and remittances to be addressed to **Entomological News, 1900 Race Street, Philadelphia 3, Pa.**

Prices per yearly volume of 10 numbers.

Private subscriptions, for personal use: in the United States, $5.00; Canada, $5.15; other countries, $5.30.

Institutional subscriptions, for libraries, laboratories, etc.: in the United States, $6.00; Canada, $6.15; other countries, $6.30.

ADVERTISEMENTS: Rate schedules available from the editor.

MANUSCRIPTS and all communications concerning same should be addressed to **R. G. Schmieder, Zoological Laboratory, University of Pennsylvania, Philadelphia 4, Pa.**

The receipt of all papers will be acknowledged and, if accepted, they will be published as soon as possible. Articles longer than eight printed pages may be published in two or more installments, unless the author is willing to pay the cost of a sufficient number of additional pages in any one issue to enable such an article to appear without division.

ILLUSTRATIONS: Authors will be charged as follows: For text-figures, the cost of engraving; for insert plates (on glossy stock), the cost of engraving plus printing. Size limit, when printed, 4×6 inches. All blocks will be sent to authors after printing.

TABLES: The cost of setting tables will be charged to authors.

SEPARATA: Members of the American Entomological Society may elect to receive, gratis, 25 offprints of their contributions. These will be "run-of-form," without removal of extraneous matter.

Those members desiring more than 25 separates, and all non-members, will receive no gratis copies. They must obtain all their separates (as reprints, with extraneous matter removed) from the printer at the prices quoted below. Authors must place their order for such separates with the editor at the time of submitting manuscripts, or when returning proof.

Copies	1–4 pp.	5–8 pp.	9–12 pp.	Covers
50	$4.35	$6.96	$10.88	$4.74
100	5.21	8.26	13.05	6.48
Add'l 100	1.74	2.60	4.33	3.48

Plates printed one side: First 50, $3.47; Additional 100's, $2.61.

Transportation charges will be extra.

ENTOMOLOGICAL NEWS

VOL. LXXII JULY, 1961 No. 7

A New Species of Stictiella from Mexico (Sphecidae: Bembicini)

JAMES E. GILLASPY, Sul Ross State College, Alpine, Texas

This description is presented to make the name available for publication of behavioral data by the collector, Dr. Howard E. Evans, for whom the species is named.

Stictiella evansi, n. sp.

Holotype Female.—Length 15 mm. Color pattern triphasal black, yellow, and clear (or hyaline), the latter limited principally to margins of metasomal sclerites. Black areas represent melanic infusion of the integument, which is otherwise clear. Yellow is developed beneath and is seen through the transparent integument. Yellow areas are as follows: pedicel and scape below; clypeus except pair of "nasal" spots; intersocketal area, not attenuated above or enclosing sockets; broad anterior orbits narrowing abruptly above; complete V above frontal pit receiving anterior ocellus; posterior orbits exceeding inner angles of compound eyes, continuous across vertex except for narrow median and lateral interruptions, not attaining the occipital suture posteriorly, and no yellow being found posterior to the occipital suture; pronotum except transverse band extending to base of each lateral lobe; mesonotum in form of nested U's, outer U based on scutellum, extending anteriorly almost to anterior margin of scutum and bordering scutum except anterior to tegulae, inner U lyre-shaped, interrupted medially; tegulae anteriorly; postscutellum except narrow anterior crescent; propodeal triangle except basal crescent and apex; postero-

lateral angles and anterior border of propodeum including spiracles; mesepisterna and metepisterna broadly above, except black along sulci; discal spot of hypoepimeral areas; coxae and trochanters apically to some extent; femora except above; tibiae except spot below on middle tibiae and except hind tibiae above; anterior tarsi; middle tarsi except distitarsi above; hind tarsi except nebulous areas above; first tergite except anterior face, invading subsutural areas basally and extending posteriorly as a broad median tongue, triradiate, narrowly connected to apical black at midline, leaving elliptic-ovate postgradular spots joined to lateral maculation; remaining tergites with elements of this pattern, but antero-median black tongue broadening on tergites 2–4 (proportionately broader on 5–6) and not connected to apical black except on tergite 5; tergite 6 without apical black; sternite 1 medially and apically; sternites 2–6 with progressively smaller lateral spots, separated by progressively broader, apically narrowing, medial black.

Vestiture inconspicuous, not concealing integumental surface, that of clypeus and anterior orbits fine, appressed, giving silvery sheen.

Head wider than thorax at posterior lobes of pronotum (1.06:1). Scape moderately stout, length about three times greatest width. Maxillae apicad of palpal base in length more than half of head width (1:1.77); maxillary palpi with six segments, labial palpi with four. Labrum longer than basal width (1.2:1). Clypeus width less than half of head width (1:2.27), narrower than distance between compound eyes at vertex (1:1.04), distinctly less than interocular distance at vertex (1:1.10); surface of clypeus slightly protuberant in lateral view, exceeding intersocketal carina; basal part of clypeus without a distinctly planate area; epistomal suture at closest point distant from antennal sockets by about one-fifth intersocketal distance (1:4.8), this at subantennal angles, between which it is slightly bowed downward; outwardly from subantennal angles sloping distinctly to tentorial angles, thence almost rectilinearly to compound eyes, where it is angulated and again almost rectilinear to lateral angles. Intersocketal carina obsolete above clypeus,

highest at midpoint of sockets, not exceeding sockets above except as weakly raised frontal line to frontal fovea, above which it is weakly impressed to anterior ocellar basin. Anterior ocellar mound circular in form, both a crest and peripherally, shallowly interrupted on midline below, deeply interrupted to lens level on midline above. Anterior ocellus a glabrous, light-pervious surface occupying the floor of the basin formed by the anterior ocellar mounds, surface delimited above by an arch-shaped sulcus, thence sloping, crescentwise, to deepest point immediately outside (below) arms of arch; arch slightly wider than long (1.14:1).

Mesosoma with punctation of mesoscutum and scutellum uniformly fine and dense. Propodeal triangle formed of rectilinear sutures, converging on posterior face of propodeum at about a 90° angle. Legs of medium build; distitarsi with scattered bristles ventrally; arolium and other median pretarsal structures not at all bulbous but padlike and scarcely evident, the claws capable of close apposition; claws all similar, uniform in curvature, outer claw of each pair very little longer. Wings beyond humeral plate two and one-half times thorax width, measured at posterior lobes (2.51:1); second cubital cell slightly narrowed above, slightly wider than high.

Metasoma with tergite 2 having smallest lateral punctures, exclusive of those in unpigmented marginal area, similar to subsutural punctation of tergite 1, tergites 3–6 with punctures progressively coarser and more sparse.

Allotype Male.—Length 16 mm. General appearance and pattern of markings fairly similar to female except more slender and maculation less extensive.

Antennae with tyloides evident on segments 4–13 (apicad only on 4) as longitudinal, broadly raised or subcarinate glabrous areas; segments 2–12 distinctly excised distally on side inward to curvature of antennae; penultimate segment without inner apical process.

Legs slender; distitarsi slender, widening apically, length more than three times greatest width, all approximately similar in form and size; anterior femora slender, only moderately thin

dorso-ventrally; anterior tarsal segments 2–4 not distinctly lobed or flattened; middle femora slender, anterior and posterior surfaces longitudinally rectilinear, parallel, posterior surface carinate-serrate, teeth relatively weak, unevenly spaced, increasing in size apically, from posterior aspect the carina and serrations rectilinear and uniformly medial with respect to dorsal and ventral surfaces except two apical teeth widely spaced, deeply divided, and diverging from line in an anterior direction; middle tibiae moderately slender, slightly exceeding femora in length when apposed; calcar of middle tibiae apically curved inward, blunt, thumblike, slender, brownish; middle basitarsi straight, slightly thickened apically, ventral surface beset with several (about 6) bristles along its length, without apical process; second and third mesotarsal segments not apically produced; posterior basitarsi unmodified.

Metasoma with seventh tergite narrowed at apex, indistinctly bilobed, dorsomedian preapical surface with well-defined, almost completely impunctate glabrous area; lateral margins above spiracular lobes inflected, groovelike, receiving dorsal margin of spiracular lobes, grooved surface bare but adjacent dorsal surface weakly produced at apex of spiracular lobes and densely set with stout, spinelike bristles; spiracular lobes moderately inflected, broadly bladelike, length only twice width, separated across venter by about one-half width of either, and with 7th sternite forming a downward arching floor to the 7th segment, opaque except very narrow membranous inner margin, at apex with broad dorsal point; surface of spiracular lobes finely lined, meshlike, glabrous except group of punctures (about 5) each with a single long hair, ventrad to spiracle; latter at upper third, slightly apicad of middle, on edge of strongly developed postspiracular pit. Sternite 2 with paired processes, sternite 6 bowed downward in apical half, somewhat keel-like anteriorly, apical margin produced to median point; eighth sternite with three terminal processes, median process turned angulately downward, thus comprising slightly less than half the measured total length of sternite, angulately margined at middle on each side; discal process not present.

Genitalia with parameres slender, ventral surface uniformly sclerotized, outwardly set with long hairs; volsellae with cuspis slender, distinctly shorter than digitus; digitus stoutly bird-head-shaped; aedeagus head slender, elongate.

This species displays various *S. pulchella* (Cresson) Group (Gillaspy, 1959) characteristics. It stands close to *S. tuberculata* (Fox) in the nature of the middle femora of the male, but has paired processes of the second sternite. The males key according to Parker (1929) to *pulchella* with only some difficulty occasioned by the weakly developed characters of the basitarsi, which are straight and only weakly emarginated throughout their length on the inner surface. In *pulchella* the basitarsi are strongly curved as well as inwardly emarginated lengthwise, and the posterior serrate carina of the mesofemora deviates dorsad almost from its origin, rather than traversing the postero-medial aspect of the femora. The female runs to Parker's couplet 37, differing from both options in having the discal marks of the scutum well developed, the mesopleura with yellow almost coextensive with black, and the sternites likewise all with both yellow and black.

Holotype female from Mazatlan, Sinaloa, MEXICO, July 18, 1959; *allotype* male from San Blas, Nayarit, Mexico, July 20, 1951, collected on sand; two *paratype* males with same data as allotype except one does not bear a "collected on sand" label; all collected by H. E. Evans. The holotype and allotype are deposited in the United States National Museum, Washington, D. C. The paratypes are in the collections of Cornell University and the author.

REFERENCES CITED

GILLASPY, J. E. 1959. A new bembicine wasp related to *Stictiella tenuicornis* (Fox), with certain phylogenetic considerations. Pan-Pac. Ent. 35: 187–194.

PARKER, J. B. 1929. A generic revision of the fossorial wasps of the tribes Stizini and Bembicini with notes and descriptions of new species. Proc. U. S. Nat. Mus. 75(5): 1–203, 15 pls.

A New Genus and Species of Firefly: Photoctus boliviae (Coleoptera; Lampyridae)

FRANK A. MCDERMOTT, Wilmington, Delaware

Occasionally one encounters a specimen which just does not fit in with accepted character combinations for classification. Such is the case with three male specimens of a lampyrid sent me from Bolivia by Sr. Luis E. Peña, of Santiago, Chile. The remarkable characters of this insect are the much reduced epipleura and an enlarged and quite certainly luminous 8th ventral segment. For this species I propose the new generic name *Photoctus,* a condensation implying the luminous 8th ventral, and the specific name *boliviae,* indicating the country of origin.

PHOTOCTUS gen. nov.

This genus is differentiated from previously described lampyrid genera by the following combination of characters:

Antennae uniramose, the rami long, flattened, contorted, hairy; antennal sockets prominent, projecting beyond the eyes.

Mandibles apically slender.

Gula membranous in forward portion.

Epipleura reduced to basal traces; no explanate, elytral margins; forward edges deflexed vertically at the humeral angles, becoming level with the disk at about midlength.

Tergites with acute lobes, directed posteriorly.

Eighth ventral segment long and broadly expanded, slightly emarginate at apex; apparently luminous.

Photoctus boliviae sp. nov.

Description of holotype:

Type locality El Palmar, Chopare, Bolivia. Collector Luis E. Peña, September 8, 1956.

Dimensions *ca.* 5.8 mm long by 2.05 mm broad; outline parallel.

Pronotum *ca.* 0.9 mm long by 1.6 mm broad at angles; nearly evenly semicircular; posterior angles acute and projecting be-

hind the middle of the sinuate base. Anterior half of margin internally deflexed, deeply so at sides of eyes; posterior half nearly flat; coarsely and densely punctate; pubescence short and scanty, principally on edges; color transparent brown. Disk convex, smooth, and shining, with scattered hairs; basal half of convex area consists of a trapezoidal brown spot not quite reaching base, with a median longitudinal channel; convexity extends forward over eyes, with a short median carina.

Scutellum dark brown, coarsely punctate, apex broadly rounded. Mesonotal plates dull dark yellow.

Elytra 4.85 mm long by 1.02 mm broad; parallel (actually somewhat spread in holotype), brown, densely rugose-punctate and markedly tricostate, the costae almost reaching the apices; no explanate margins and only basal traces of epipleura; the outer edges deflected vertically at the humeral angles and gradually becoming level with the disk at about midlength. Fairly dense, short, oblique pubescence; no secondary pubescence observable.

Tergites all very dark brown, 4th to 7th with posteriorly directed acute lobes. Pygidium black, trilobed, wider and longer than the 8th ventral.

Prosternum dull yellow, mesosternum yellowish brown, metasternum cloudy reddish brown.

Ventral segments 2 to 5 dark brown, 6th with posterior edge pale, and 7th almost entirely so, both broadly emarginate; pubescence short and appressed. 8th ventral as long as 6th and 7th combined, broadly expanded, nearly as wide as 7th and narrowly emarginate in middle of posterior edge; median longitudinal sulcus; pale salmon color.

Legs short; fore and intermediate light brown, posterior darker; tibial spurs absent. First posterior tarsal article as long as the next three combined; 4th with very small pulvilli; 5th nearly as long as first. Claws small, sharp, simple.

Head deeply set in "collar" of the prothorax; frons concave, very dark brown, rugose; interocular margins not divergent; 0.9 mm across eyes and 0.5 mm between them above antennal sockets; eyes large and contiguous below.

Mandibles very small, pale; distal portion slender and very sharp, approaching Green's modified type. Terminal article of maxillary palpi conical and but little thicker than the first three articles; labial palpi very small, terminal crescentic.

Gula membranous in the small forward portion which is visible.

Antennae 2.3 mm long, articles 3 to 10 dark brown, subequal in length, and each bearing a single pale, flattened, hairy ramus 3 to 5 times as long as the article from which it arises, narrowly reniform; rami contorted, not folded fanwise. 11th article similar to the ramus on the 10th. Antennal sockets and mouth parts project forward beyond the eyes.

Abdominal spiracles not discernible; presumably in the pleural fold.

Aedeagus not extracted, but protruded in one paratype, showing a compact conical form with the median lobe visible between the lateral lobes and curved upwards, the tips of all three approximate; two small lateral knobs.

Female unknown.

Holotype is being deposited in the U. S. National Museum as No. 65674; one *paratype* in the California Academy of Sciences, and one in my collection. The paratypes are slightly smaller than the holotype, and one is somewhat less strongly pigmented; the 6th ventral may be entirely brown.

It would be very interesting to find the female of this species; probably larviform and brightly luminous. The luminous conduct of the male was not reported.

Concerning the Neogeophilidae, with Proposal of a New Genus.[1] (Chilopoda: Geophilomorpha: Neogeophilidae)

By R. E. Crabill, Jr., U. S. National Museum, Smithsonian Institution, Washington, D. C.

(*Continued from p. 159*)

Cryptostrigla silvestri, new species

Holotype: ♀. Guatemala: Department of Alta Verapaz, Semococh (according to O. F. Cook's note, about 48 km. southeast of Coban). G. P. Goll, leg. U. S. National Museum catalogue of myriapod types: 2606; chilopod type C-147; see slides StC: 76 and 77.

Introductory. Length, about 32 mm. Pedal segments, 69. Body shape: Essentially of uniform width, the final 4–5 segments slightly attenuate. Color: Considerably discolored from long immersion in alcohol, hence uniformly sordid light brown.

Antennae. Each is broken; left with 5 articles, right with 4 articles. Each basal article much wider than long, the remaining articles approximately as long as greatest width. Vestiture evidently becoming suddenly denser on the fifth article. Cephalic Plate. Greatest length, 0.544 mm; greatest width, 0.579 mm, thus somewhat wider than long. Shape: Anteriorly broadly pointed, the two sides meeting to form an obtuse angle; laterally strongly, evenly excurved; posterior margin essentially straight. Areolation coarse. Without frontal or other sutures or sulci. Setae short and sparsely disposed. A narrow, central portion of the prebasal plate is exposed. Clypeus (Fig. 6). Paraclypeal sutures nearly straight; complete (not curving postero-laterally as in Silvestri's figure of *primus*). Setae, few, as shown; prelabral setal pair absent. Clypeal areas and plagulae absent; areolation coarse and essentially uniform. Each bucca well-defined by strong sutures; each with a weak transbuccal suture; anteriorly with a few small setae. Labrum (Fig. 6). Consisting of one obscure, weak, hyaline, undivided piece

that projects posteriorly in a gentle convex arc; labral teeth very short and delicate, hyaline. Labrum continuous on each side with a delicately sclerotized bar (part of the clypeus) that meets each labral fultura. MANDIBLE (Fig. 1). Shaft relatively short; body of the mandible relatively long and massive; distally with a row of simple hyaline teeth, these very long and flat, rather blunt. FIRST MAXILLAE (Fig. 2). Coxosternum completely divided medially into right and left halves; lappets absent. Each coxosternal half surmounted by a broad, lobelike structure (which may represent a highly modified telopodite); each lobelike structure apically with an indistinct membranous area but otherwise without sutures, divisions, etc.; lappets absent. SECOND MAXILLAE (Fig. 2). Coxosternum medially rather narrow, neither divided nor suturate midlongitudinally; postero-laterally somewhat extended; entire posterior margin weakly areolate and regionally membranous; metameric pore openings conspicuous. Telopodite consisting of three distinct articles; basal article entirely without dorsal and ventral condyles; third article rather ovate in outline; apical claw short and broad, pointed, not excavate, anterior edge smooth but posterior edge with about 3 sharp teeth, thus giving claw superficially a bifid appearance (Fig. 8).

PROSTERNUM (Fig. 3). Very broad. Pleuroprosternal sutures complete, terminating dorsolaterally. Abortive subcondylic sclerotic lines present, these continuous with the pleuroprosternal sutures posteriorly but not passing to or toward their respective prehensorial condyles. Anterocentrally with a pair of obscure but well-sclerotized and rounded denticles. PREHENSORS (Fig. 3). When flexed, falling far short of anterior cephalic margin. Denticles absent on all articles. Ungula long and extraordinarily straight, falciform; dorso-ventrally very flat, bladelike; posterior edge finely dissected to form about 6 tiny, irregular serrations. Poison calyx located at upper end of trochanteroprefemur, cordiform in shape; its duct passing along anterior edge of ungula to open far short of apex. Poison gland very long, passing out of trochanteroprefemur, apparently extending posteriorly nearly as far as 1st pedal segment.

TERGITES. Basal plate anteriorly weakly concave to reveal a small portion of prebasal plate centrally. Remaining tergites coarsely areolate, very sparsely setose; without sulci. PLEURITES. Agreeing closely with those of *E. mexicanus* (Silvestri's Figs. 1–5, p. 357). Paratergites absent. Spiracles on anterior third of body weakly horizontally elliptical, thereafter becoming rounder. LEGS (except ultimates). First legs only slightly shorter and thinner than those following. All legs short and notably robust, not becoming longer or less robust posteriorly on body. Setae short and sparse but more numerous than on tergites and sternites. Pretarsi (Fig. 4): Each fundamentally consisting of a rather bulbous base and a prominent claw proper;



Cryptostrigla silvestri, new species, holotype.

1. Left mandible (inner surface).

claws proper from the 1st through the 33rd each with a conspicuous antero-ventral, ventrally directed tooth, tooth of 1st pretarsus small, thereafter ventral teeth increasing in size, becoming smaller again on the 31st and 32nd pretarsi; each tooth-bearing pretarsus with minute serrations on the ventral edge of claw proper; each pretarsus (1–68) with two basal accessory spines, the posterior always minute, the anterior very robust and long on those pretarsi with ventral teeth (on 1 through 33), thereafter becoming smaller and thinner. STERNITES. Sulci, sutures, carpophagus-structures, porefields, depressions, metasternite projections all absent. Setae short and sparse. Areolation weak. On anterior third of body the intersternites are weakly divided midlongitudinally; on posterior two-thirds of



EXPLANATION OF FIGURES

Cryptostrigla silvestri, new species, holotype.

2. First and second maxillae (ventral aspect, setae deleted).
3. Prosternum and right prehensor (ventral aspect).
4. Pretarsus and tarsus of 14th left leg (posterior surface, all setae shown). a, minute serrulations on plantar edge of claw proper. b, ventral tooth of claw proper. c, hypertrophic anterior accessory spine. d,

body intersternites become wider anteroposteriorly, more band-like; undivided centrally.

ULTIMAE PEDAL SEGMENT (Figs. 5, 7). Pretergite intimately fused with tergite proper; the suture separating them is persistent but vestigial and weak; pretergite evidently without pleurites. Tergite: Greatest length exceeds greatest width; sides nearly parallel, weakly convergent; posterior margin medially extended to form a blunt point, the two sides (of the rear margin) thus forming an obtuse angle. Presternite intimately fused with sternite, the vestigial suture separating them is present but discernible with difficulty. Sternite: Sides weakly convergent; the *true* posterior margin medially very deeply excavate, the two corners extended posteriorly in long sharp points; the sternite intimately fused with the pregenital sternite, the intervening suture barely discernible but persistent. Coxopleuron: Moderately inflated; dorsally, laterally and ventrally with small, irregularly disposed freely-opening pores; without porepits of parasternital fossae; setae short and very sparse. Leg: Notably longer and thinner than penult leg; with 6 articles distal to coxopleuron; setae short and somewhat more numerous than on other legs; tarsus consisting of two articles, the distotarsus slightly longer than the proximotarsus; pretarsus represented by a minute sclerotic point (seen only at 645 ×), hence an unguiform or tuberculate pretarsus is absent.

(in dashes) depressor tendon of the pretarsus. e, claw proper of the pretarsus. f, minute, atrophied posterior accessory spine. g, (in dashes) condyle of pretarsus.

5. Ultimate pedal segment and postpedal segments (ventral, setae deleted). a, penultimate pedal sternite. b, presternite of ultimate pedal segment. c, ultimate pedal sternite. d, vestigial, extremely obscure but persistent suture separating the true ultimate pedal sternite (c) and the pregenital sternite (e). e, pregenital sternite. f, genital sternite. g, (in dashes) concealed terminal pore. h, gonopod.

6. Clypeus, labrum, and right and left buccae (ventral aspect, all setae shown).

7. Rear body segments (dorsal aspect, setae deleted). a, pretergite of penultimate pedal segment. b, tergite of penultimate pedal segment. c, last spiracle of left side. d, pretergite of ultimate pedal segment. e, obscure transverse suture separating pretergite (d) and tergite (f). f, tergite of ultimate pedal segment.

8. Third article and claw of 2nd maxillary right telopodite (ventral aspect, all setae shown).

Postpedal Segments (Fig. 5). Pregenital sternite anteroposteriorly very long, passing forward to and fusing intimately with the ultimate pedal sternite from which it is separated by a vestigial, obscure suture, thereby causing the last pedal sternite to appear much longer than it actually is. Each gonopod is conical, long; entirely without a discernible interarticular suture or other indication of division, hence is secondarily uniarticulate. Terminal pores present, small, concealed.

On the Rank and Possible Affinities of the Neogeophilidae.—The real importance of the discovery of this specimen derives from the unusual opportunity it affords for the direct examination of a member of this peculiar and systematically unsettled group. Careful examination of the animal testifies to the thoroughness of Silvestri's morphological diagnosis; as we have seen, the accuracy of his report is questioned pertinent only to four points, none of which would alter Silvestri's original contention that the group is suprageneric in rank. However, my examination of the single specimen, together with a careful reconsideration of Silvestri's published data, at this time do not permit any other *confident* conclusion than his own, that the rank of the group is probably suprageneric. For reasons explained below it seems preferable for the time-being to regard the neogeophilids as members of a distinct, aberrant family within what I shall call the geophilid-sogonid-gonibregmatid complex of families.

It seems clear that, while belonging to this family complex, the neogeophilids appear to be referable to no one of these families, at least as they are currently defined. At the same time, many of the neogeophilid structures *individually* are reminiscent often of closely similar counterparts that are discernible within this great suprafamilial section of the Geophilomorpha.

The problem of determining the rank and affinities of the Neogeophilidae is by no means reducible merely to one of deciding which is the best and most reasonable of several alternatives in the light body of well-understood and digested morphological information whose details are familiar to everyone. On the contrary, the interpretive problem is necessarily super-

imposed and dependent upon a much more formidable one in this case: many of the most critical facts upon which our inductions must depend are actually representative of a persistent legacy of deficient information which is further complicated by frequent breakdowns in interpersonal understanding. First, there are huge gaps in our knowledge of the full spectrum of the Geophilomorpha: certainly, many groups and species still await discovery. Secondly, in the case of the majority of recognized groups and species we must remain ignorant of the nature, or even of the existence, of many critical diagnostic features if, as is often unavoidable, we must depend for full, precise information upon published descriptions. Finally, it is not infrequently true that even when critical features are treated, their explication is so loose and imprecise, so subjective and cryptic, or even so faulty as to preclude the reader's gaining an accurate or sufficiently detailed understanding of them.

These several difficulties create a particular problem for the categorical assessment and group assignment of the neogeophilids whose conceivable closest relatives as groups are themselves often systematically unsettled, descriptively obscure, and evidently poorly known in terms of the species and supraspecific groups that exist but are undiscovered. The particular problem that is posed is how to interpret the structures about which we believe we have reasonably accurate, meaningful information under these circumstances. Specifically, are these presumably homeomorphic structures evolutionarily conservative, being derivative from a single and immediate preexisting source, or are some or all of them convergent and polyphyletic, having been derived independently, compelled alike by adaptive pressures in separately evolving, remotely-related lines?

Under the circumstances, and with reference to the Neogeophilidae, it seems impossible to settle this question now. We do not know enough about a sufficient number of structures and structural complexes. We do not know enough about the geophilidiform centipedes to be able to distinguish between convergencies and immediately derived structures and forms. At the same time, it is desirable to make mention of some homeo-

morphic characters that eventually may or may not prove to signify close phylogenetic linkages between the neogeophilids and certain other geophilidiform groups.

In general body habitus the Neogeophilidae bear an undeniable resemblance to the Sogonidae and some resemblance to the Dignathodontidae, although their overall similarity, e.g., in head and body shape, to the latter may well be only superficial and adaptively convergent.

A rather homogeneous, poorly-known assemblage of geophilid-like centipedes, the sogonids, apparently are restricted to the more northern New World tropics and to adjacent parts of North America where they are evidently incursive from the south. Established as a family and almost entirely described by Professor Chamberlin, the Sogonidae are clearly abundant in the neotropics where many new groups and species probably await discovery. Like the neogeophilids, they are all small, delicate creatures. Tiny short heads, delicate prehensors, simple and apparently vestigial labra,[2] simple mandibles and, reportedly in some sogonids, aberrant maxillary configurations suggest a general similarity whose explanation on the grounds of immediate evolutionary derivation, however, can hardly be very convincing in our present state of knowledge. Nonetheless, while differing in several critical features, the two groups, as

[2] The sogonid labrum has been inaccurately described repeatedly. Originally Chamberlin described it as being ". . . of a single piece apparently free laterally but fused in the middle; . . ." (1912, p. 432). Completely misinterpreting Professor Chamberlin's statement, Attems wrote the following in his key to families (1929, p. 27): "Oberlippe aus einem ungeteilten Stück bestehend." The first description is cryptic, the second erroneous. On the one hand, they are suggestive of the single, or unipartite, type of labrum, such as that labral type that is characteristic of the schendylids or himantariids. On the other hand, they fail to stress what is really significant, that the sogonid labrum is fundamentally of the tripartite geophilid type, departing from it in the apparent direction of degeneracy. In *Sogona minima* there are two prominent sidepieces which are relatively well-developed, discrete, and widely-separated. Most importantly, the midpiece has either atrophied entirely, or else it has fused imperceptibly with the broadly intruded midclypeal extension. In summary: We can only describe the sogonid labrum as being fundamentally tripartite and lacking a *distinguishable* midpiece.

we know them of course, do appear superficially to be rather similar. It is difficult to ignore the possibility, however remote, that the neogeophilids and sogonids may represent closely-related, aberrant evolutionary experiments that were fragmented together from some ancient geophilid stock. Similarly, the dignathodontids and aphilodontine geophilids perhaps represent separate and now nominate variations upon an original, basic geophilid theme.

The first maxillary coxosternum of the neogeophilids, being totally divided into right and left halves, is curiously suggestive of its homologue in *Himantosoma,* a genus that Attems placed in Gonibregmatidae but which Verhoeff regarded as the basis of a separate family. Again, whether these divided coxosterna are merely convergent or are evolutionarily derivative from a common precursor is impossible to determine now. It may, however, be significant that *Himantosoma* lacks the bizarre anterior maxillary lobes that signalize all known neogeophilids.

As I have noted above, the neogeophilid labrum appears by direct inspection to be simple and degenerate and reminiscent of that of the sogonids, inasmuch as that is also evidently hypertrophied. This is not to imply that they resemble each other very closely; they do not. At the same time, essentially the neogeophilid type of labrum may be seen in certain Gonibregmatidae. For that matter, the same labral type is found in certain ballophiline Schendylidae, which do not seem very closely related to the whole section of the Geophilomorpha here under discussion. Without much doubt, quite similar, if not occasionally identical, labra have arisen *independently* at least in some unrelated geophilomorphs.

The neogeophilid mandible, equipped only with a simple row of delicate hyaline and homogeneous teeth, apparently can tell us little, except that the neogeophilids *may be* more closely related to the geophilid-dignathodontid-gonibregmatid-sogonid complex than to any other. But even in this regard we are hardly entitled to conclude with an emphatic finality that this simple type of mandible *in every instance* can only be indicative of monophyletic origins. It is by no means impossi-

ble that, let us say, through convergency, or by whatever mechanism, the simple geophilidiform mandibles was developed independently in remotely related geophilomorphs. If we ignore the venerable mandibular criterion momentarily, then several rather striking structural similarities existing between the neogeophilids and oryids could possibly take on new significance.[3]

The neogeophilid pretarsi are evidently unique. The extraordinary ventral teeth, hypertrophic anterior accessory spines, and serrulate plantar edges must function as a unit to facilitate traction upon or adherence to the surfaces over which their possessors move. Analogous, though evidently not wholly homologous, adaptive devices are known to occur in some other geophilomorphs. For instance, the gonibregmatid genus *Eucratonyx,* while lacking a ventral pretarsal tooth, has a conspicuously introrse claw proper which, in conjuction with a hypertrophic anterior accessory spine, probably affords a firmer foothold for locomotion over rough surfaces or for stationary clinging. Again, massive development of the claw proper and of one or both of its accessory spines has been noted in certain schendylids (e.g., *Pectiniunguis*). It seems quite likely that cryptophiles such as these geophilomorphs would be inclined to evolve efficient hold-fast devices independently: their existence depends upon adaptation to a variety of crevice-cranny habitats wherein, one would think, adaptive pressures would place a premium upon the ability to squeeze through tight, narrow confines and to anchor firmly against forceful removal by predators.

Summing up: The structures that signalize the known Neogeophilidae tell us little conclusively about their interfamilial affinities. Many of these structures could very well represent adaptive convergencies that obscure rather than illuminate the ranks and affinities of groups. While most individual neogeophilid features have often quite similar counterparts in various other geophilomorph groups, in no case is there a concordance of structural identities that could justify an unequivocal statement

[3] By the same token, if we ignore the mandibular criterion, a number of features in certain oryids suggest possibilities that have not received serious consideration.

of close phylogenetic affinity at this time. It is conceivable that the neogeophilids and sogonids could reflect a community of descent, although, admittedly, this is a highly speculative suggestion for which present evidence is limited and frankly unconvincing.

A Review of the Neogeophilidae

Neogeophilidae

Neogeophilinae, Silvestri, 1918, p. 352.
Neogeophilidae, Attems, 1926, p. 365.

Distinguishing Criteria.—1st maxillary coxosternum completely divided into right and left halves, each half surmounted by a large, uniarticulate lobe. Pretarsi of the more anterior legs each ventrally with a prominent tooth that is continuous with the claw proper. First article of second maxillary telopodite basally without condyles.

Extended Characterization.—Antennae slightly attenuate distally. Cephalic plate very slightly wider than long to slightly longer than wide; frontal suture absent. Prebasal plate at least slightly exposed. Clypeus with complete paraclypeal sutures; without clypeal areas or plagulae. Labrum comprising a delicate undivided bar, wholly amalgamated with postero-central clypeus, with delicate hyaline teeth, these long, flat, rather blunt; proximal shaft relatively short when compared with the longer, heavier distal dentigerous portion. First maxillae: Coxosternum medially completely divided, the right and left sides thus entirely discrete; each coxosternal half with a large lobate structure in place of the usual structures; lappets absent. Second maxillae: Coxosternum medially undivided, not suturate; teleopodite basal articles without discernible condyles; apical claw with a few delicate spiniform projections arising from posterior edge. Prosternum with complete pleuroprosternal sutures; with or without complete subcondylic sclerotic lines. Prehensors: When closed falling far short of anterior cephalic margin; articles without denticles; ungulae long and falcate, flattened dorso-ventrally, posterior edge finely, irregularly serrulate.

Tergites not sulcate. Paratergites absent. Legs robust and short. Pretarsi of the more anterior legs each with a conspicuous ventral tooth and equally conspicuous hypertrophic anterior accessory spine. Sternites not sulcate; without porefields or carpophagus-structures; the more posterior intersternites broadly bandlike and not suturate midlongitudinally. Ultimate pedal segment: Pretergite either separated from its tergite by a distinct transverse suture, or, if fused intimately with tergite, separated from it by an obscure suture or else apparently without an intervening suture. Coxopleuron with freely opening pores, without dorsal or ventral porepits or porigerous fossae; moderately inflated. Sternite either distinguishable from or intimately fused with the pregenital sternite; ultimate tarsus uniarticulate in the males, biarticulate in the females; pretarsus essentially absent. Terminal pores present but concealed. Each gonopod biarticulate (with persistent intervening suture), or uniarticulate (without intervening suture).

Range: Known only from Mexico and Guatemala.

Known from three monotypic genera, as follows.

NEOGEOPHILUS Silvestri

Neogeophilus Silvestri, 1917, p. 352.

Type-species: *Neogeophilus primus* Silvestri, 1917. (Original designation and monotypic.)

Diagnosis.—With the characters of the family, of which the following signal characters are distinctive. (1) Head slightly longer than wide. (2) Paraclypeal sutures apparently arching outward, apparently not terminating on the rear clypeal margin (see discussion under foregoing Notes). (3) Prosternal denticles absent. (4) Prosternal subcondylic sclerotic lines passing to and esesntially meeting their respective condyles. (5) Ultimate pedal pretergite and tergite, and sternite and pregenital sternite separated by distinct sutures. (6) Female gonopods biarticulate.

Inclusive species: Known only from *N. primus* Silvestri: with the characters of the genus, in addition ♂ with 81 pedal

segments, 34 mm. long; only known and type locality, Cuernava, State of Morelos, Mexico.

EVALLOGEOPHILUS Silvestri

Evallogeophilus Silvestri, 1917, p. 355.

Type-species: *Evallogeophilus mexicanus* Silvestri, 1917. (Original designation and monotypic.)

Diagnosis.—With the characters of the family, of which the following signal characters are distinctive. (1) Head considerably longer than wide. (2) Paraclypeal sutures terminating posteriorly on the rear clypeal margin. (3) Prosternal denticles present. (4) Prosternal subcondylic sclerotic lines passing to and essentially meeting their respective condyles. (5) Ultimate pretergite and tergite, and sternite and pregenital sternite intimately fused, apparently without intervening sutures. (6) Female gonopods biarticulate.

Inclusive species: Known only from *E. mexicanus* Silvestri: with the characters of the genus, in addition ♂ with 63, ♀ with 67 pedal segments; to 30 mm. long; only known and type locality, "Jalapa" (in full, Jalapa Enriquez), State of Veracruz, Mexico.

CRYPTOSTRIGLA, new genus

Type-species: *Cryptostrigla silvestri,* new species. (Present designation and monotypic.)

Diagnosis.—With the characters of the family, of which the following signal characters are distinctive. (1) Head somewhat wider than long. (2) Paraclypeal sutures terminating on clypeal margin. (3) Prosternal denticles present. (4) Prosternal subcondylic sclerotic lines abortive, not passing across prosternal corner to or toward their respective condyles. (5) Ultimate pretergite and tergite, and sternite and pregenital sternite intimately fused, the intervening sutures still discernible. (6) Each female gonopod single, the two articles having fused without trace of intervening suture.

Inclusive species: Known only from *C. silvestri,* new species:

with the characters of the genus; in addition ♀ with 69 pedal segments, 32 mm. long.; only known and type locality, Semococh, Department of Alta Verapaz, Guatemala.

References

Attems, C. Graf von. 1926. Kükenthal-Krumbachs Handbuch der Zoologie 4 (3+4): 241–402. 1929. Geophilomorpha, in Das Tierreich, Lief. 52: 1–388.

Chamberlin, R. V. 1912. The Geophiloidea of the South-eastern States. Bull. Mus. Comp. Zool. Harvard 54: 405–436.

Silvestri, F. 1918. Descrizione di due nuovi generi di Geophilidae (Chilopoda) del Messico. Boll. Laborat. Zool. Gener. Agrar. Portici 12: 352–360.

Supplementary Records of Meloid Beetles (Coleoptera) of the West Indies

By Richard B. Selander and John K. Bouseman,
Department of Entomology, University of
Illinois, Urbana

Since the completion of our report on the Meloidae of the West Indies (Selander and Bouseman, 1960, Proc. U. S. Nat. Mus., vol. 111, pp. 197–226), we have received from Patricia and Charles Vaurie, American Museum of Natural History (AMNH), a series of specimens collected by them in 1960 on the islands of Guadeloupe, Jamaica, and Martinique and from M. W. Sanderson, Illinois Natural History Survey (INHS), a series of specimens collected by him in 1959 on Cuba. A few other specimens of West Indian meloids have also come to our attention. In publishing the records of this supplementary material, we again take pleasure in acknowledging the cooperation of our colleagues.

In order to avoid repetition, we will list here the localities and dates for the Vaurie material. Guadeloupe: Domaine Duclos, 600 ft., June 24–28 and July 7; Les Saintes, Terre de Haut,

July 2–5; and Matouba, 1900 ft., June 29. Jamaica: Hardware Gap, 4800 ft., July 13–15. Martinique: Absalon, rain forest, June 16; Diamant, June 17; and Morne Rouge, June 13.

Cissites maculata (Swederus). GUADELOUPE: Domaine Duclos, AMNH, 2; Les Saintes, Terre de Haut, AMNH, 1. HISPANIOLA: Sánchez, Dominican Republic, June 3–12, 1915 and February 1916, AMNH, 3.

Pseudozonitis marginata (Fabricius). GUADELOUPE: Domaine Duclos, AMNH, 64; Matouba, AMNH, 3. JAMAICA: Hardware Gap, AMNH, 1. MARTINIQUE: Absalon, AMNH, 5; Diamant, AMNH, 3; Morne Rouge, AMNH, 3. PUERTO RICO: Mayagüez, May 1924, Cornell University, 1.

The specimens from Guadeloupe are assignable to our color class 0 and those from Jamaica and Puerto Rico to class 2. In the series from Martinique, two specimens are representative of class 0, one of class 2, two of class 3, and six of class 4. One of the specimens of the last class has the pale elytral vitta reduced to half its usual width. In a few of the specimens from Guadeloupe and in one from Martinique the head is fuscous. Females outnumber males in the new material four to one.

Pseudozonitis obscuricornis (Chevrolat). GUADELOUPE: Domaine Duclos, AMNH, 12; Matouba, AMNH, 1. JAMAICA: Kingston, 1958, M. Locke, Selander collection, 1.

The specimen from Jamaica is heavily marked. Those from Guadeloupe have the elytral vitta either poorly developed or absent.

Nemognatha punctulata LeConte. CUBA: Lower slopes of Loma (Pico) del Gato, Sierra Maestra, Oriente Province, May 26, 1959, on Compositae, M. W. Sanderson, INHS, 9; between Santa Lucia and Nuevitas, Camaguey Province, June 8, 1959, M. W. Sanderson, INHS, 1.

An annotation should be made to couplet 6 of our key indicating that the color of antennal segment I is variable in this species. In all the specimens that we studied earlier this segment is yellow. However, in three of the specimens recorded above it is piceous and in four it is completely black.

Melanagromyza tiliae (Coud.) (Diptera: Agromyzidae) Reared from Linden Bark Galls at London, Ontario

By W. W. Judd, Department of Zoology, University of Western Ontario, London, Ontario

On March 18, 1957, six twig galls were collected from basswood trees, *Tilia americana*, on the campus of the University of Western Ontario at London, Ontario. They were identified as Linden Bark Galls by using keys and descriptions in Felt (1940). Each gall was a smooth swelling along the side of the twig and was one-half an inch long, three-sixteenths of an inch wide and bulged one-eighth of an inch above the surface of the twig. The bark covering a gall was as smooth as the bark on the surrounding twig and was of the same color. One gall was dissected and was found to be composed internally of soft wood. The remaining galls were kept in the laboratory in a corked vial. On April 9 a fly emerged from one of the galls through a circular hole 1.5 mm. in diameter. On April 10 the other four galls were dissected and yielded one brown puparium each. The fly was identified as *Melanagromyza tiliae* (Coud.) by Mr. G. E. Shewell, Entomology Research Institute, Department of Agriculture, Ottawa. This species was described by Couden (1908) from specimens reared from twigs of basswood and its taxonomic relationships have recently been discussed by Shewell (1953) and Frick (1957).

On September 7, 1957, seven more galls were collected from basswood on the university campus and a few days later three more adult *M. tiliae* emerged from them. The four specimens reared at London are deposited in the collection of the Department of Zoology, University of Western Ontario. Felt (1940) records that this fly produces its gall in summer. The specimens from which the species was described by Couden were reared from galls in Missouri in April (Couden, 1908; Frick, 1957), and the first fly reared at London emerged in the laboratory in this month. It is thus evident that this insect habitually overwinters in the gall on the twig.

References

Couden, F. D. 1908. Proc. Entomol. Soc. Washington 9: 34–36.
Felt, E. P. 1940. Plant galls and gall makers. Comstock Publ. Co. Ithaca.
Frick, K. E. 1957. Ann. Entomol. Soc. Amer. 50: 198–205.
Shewell, G. E. 1953. Can. Entomol. 85: 462–470.

Psectra diptera (Burmeister) in Wisconsin (Neuroptera: Hemerobiidae)

By Alvin L. Throne, University of Wisconsin, Milwaukee

Psectra diptera (Burmeister) has been recorded from Austria, Denmark, England, Finland, Germany, Holland, Ireland, Italy, Scotland, Siberia, Sweden, Switzerland and the Ukraine (Killington, 1936). It has been recorded from the northeastern United States, as far west as Michigan and as far south as Virginia and West Virginia, and from Ontario (Carpenter, 1940). It has also been recorded, as *Hemerobius delicatulus* Fitch, from Illinois (Hagen, 1861). Banks, 1905, refers to specimens from Ithaca, New York, the Agricultural College, Michigan, and Franconia, New Hampshire. He also mentions Fitch's specimens collected in northern Illinois in October. Recently new records have been reported from Connecticut, Maryland and Virginia (Mac Leod, 1960).

In my extensive collecting of Neuroptera in Wisconsin, supported by the Research Committee of the University of Wisconsin on funds from the Wisconsin Alumni Research Foundation, I have obtained four specimens of *Psectra diptera*. All of the specimens I collected are macropterous. They were all taken in my backyard in Shorewood, a residential suburb of Milwaukee.

It is interesting to note that all four of my specimens were taken in a light trap. The trap was hung from a clothes line with the top of the funnel approximately three and a half feet from the ground. Killington suggests that both micropterous

and macropterous forms are most likely unable to fly. Mac Leod gives evidence that perhaps some individuals are able to fly. The position of my trap with reference to the ground, building and taller plants and the nature of the trap itself strongly suggest that macropterous individuals can fly.

Carpenter (*op. cit.*) gives the average wing length as 6 millimeters. Killington (*op. cit.*) gives total wing expanse as about 7 mm. and reports Banks as giving a wing expanse of 5–6 mm. Mac Leod (*op. cit.*) in measuring ten Nearctic specimens found the average length of forewing as 3.82 ± 0.25 mm. and total wing expanse as 8.57 ± 0.54 mm.

The following wing measurements of the specimens I collected were made with an ocular micrometer.

Date Collected	9-VIII-1957	9-VIII-1957	27-VIII-1958	2-VIII-1959
Length of forewing	5.08 mm.	3.85 mm.	4.00 mm.	4.61 mm.
Body width at mesothorax	1.08	0.77	0.77	0.92
Total wing expanse	11.24	8.47	8.77	10.14

LITERATURE CITED

BANKS, N. 1905. Trans. Am. Ent. Soc. 32: 21–51.
CARPENTER, F. M. 1940. Proc. Amer. Acad. Arts Sci. 74: 193–280.
HAGEN, H. 1861. Smithsonian Inst. Miscell. Coll.: Washington, D. C.
KILLINGTON, F. J. 1936. A monograph of the British Neuroptera. Volume I. Ray Soc.: London.
MAC LEOD, E. G. 1960. Entom. News 71: 231–236.

Review

A MANUAL OF COMMON BEETLES OF EASTERN NORTH AMERICA. By Dillon, E. S., and L. S. Dillon. Pp. viii + 884; 554 text figures and 85 plates (4 colored). Row, Peterson and Co., Evanston, Illinois, 1961. Price, $9.25.

Here, at long last, is a really workable manual of beetles, and one that seems to have just about every imaginable virtue.

Heretofore Blatchley's was the standby, but, using it, the inexperienced often ended with more incorrect than correct determinations. In this book, the Dillons have selected the 1200 commonest beetles in 64 families, and keyed them carefully to families, genera, and species. The diagnostic features are illustrated by 554 text figures, and there is a habitus picture and a description of each species. The species selected, it is believed, include about 90% of all beetles commonly taken in the region; for most of the remainder the bibliography, arranged by families, will lead one to recent revisions and synopses.

The very form and appearance of this manual add to the pleasure of using it. It is not too bulky (page size, $5\frac{1}{2}'' \times 8\frac{1}{4}''$) and is neatly bound in semi-flexible cloth with rounded corners. It should make many new friends for the Coleoptera, including also hobbyists who will find delight in collecting beetles and seeking out their names.—R. G. SCHMIEDER.

Corrodentia in Cliff Swallow Nests

By WILLIAM F. RAPP, JR., Nebraska State Department of Health, Lincoln, Nebraska

A number of Corrodentia were obtained recently from cliff swallow nests (*Petrochelidon pyrrhonota pyrrhonota*) by means of a Berlese funnel. These nests had been collected from the Nebraska end of the Yankton bridge (South Yankton, Cedar County, Nebraska) on August 11, 1955.

The specimens were submitted to Dr. Kathryn B. Sommerman who determined them as belonging to the *Liposcelis bostrychophilus* complex, family Liposcelidae.

It is interesting to note that Hicks (1959)[1] does not list *L. bostrychophilus* as occurring in the nests of the cliff swallow. However, *L. divinatorius* has been reported as occurring in the nests of other swallows of the Family Hirundinidae.

[1] HICKS, E. A. 1959. Occurrence of insects in birds' nests. Iowa State College Press.

Entomologist's Market Place

ADVERTISEMENTS AND EXCHANGES

Advertisements of goods or services for sale are accepted at $1.00 per line, payable in advance to the editor.

Notices of wants and exchanges not exceeding three lines are free to subscribers.

All insertions are continued from month to month, the new ones are added at the end of the column, and, when necessary, the older ones at the top are discontinued.

Butterflies. Wish to exchange specimens for Japanese species. Please write to Ichiro Nakamura (Boy, age 16), 26 Aza-Nichiyama Obayashi Takarazuka-shi, Hyogo-Ken, Japan.

Phasmidae of nearctic area desired alive. Purchase or trade, drawing on large stock of major orders, worldwide. Domminck J. Pirone, Dept. Entomology, Cornell University, Ithaca, N. Y.

Nitidulidae and Rhizophagidae wanted in exchange for European beetles of all families. O. Marek, Zămberk 797, Czechoslovakia.

Wanted and Needed. We are compiling a history of entomology, and particularly, at present, of the amateur insect clubs that flourished 50 to 75 years ago. Will you who have knowledge of such early clubs or societies advise me, giving facts on the time of existence, members, etc., which you may have. J. J. Davis, Dept. of Entomology, Purdue University, Lafayette, Indiana.

Cockroaches (Blattoidea) of Japan, Okinawa, Formosa (Taiwan), and the Philippines are being studied in cooperation with Dr. K. Princis. Loans of specimens from that area are desired. A. B. Gurney, U. S. National Museum, Washington 25, D. C.

Orthoptera. Gryllinae (except domestic sp.) and **Pyrgomorphinae** of the world wanted in any quantity for work in morphology, taxonomy, cytology, and experimental biology; dry, or in fluid, or living. Write D. K. Kevan and R. S. Bigelow, Dept. of Entomology, McGill University, Macdonald College, Quebec, Canada.

Beetles of the world wanted, all species in exchange for American beetles, moths and butterflies. James K. Lawton (age 18), 7118 Grand Parkway, Wauwatosa 13, Wisconsin.

Used genuine Schmidt boxes, excellent condition, at less than half price. H. W. Allen, Box 150, Moorestown, N. J.

Important Mosquito Works

MOSQUITO ATLAS. Part I. The Nearctic Anopheles, important malarial vectors of the Americas, and Aedes aegypti and Culex quinquefasciata

MOSQUITO ATLAS. Part II. The more important malaria vectors of the Old World: Europe, Asia, Africa and South Pacific region

By Edward S. Ross and H. Radclyffe Roberts

Price, 60 cents each (U. S. Currency) with order, postpaid within the United States; 65 cents, foreign.

KEYS TO THE ANOPHELINE MOSQUITOES OF THE WORLD

With notes on their Identification, Distribution, Biology and Relation to Malaria. By Paul F. Russell, Lloyd E. Rozeboom and Alan Stone

Mailed on receipt of price, $2.00 U. S. Currency. Foreign Delivery $2.10.

For sale by the American Entomological Society, 1900 Race Street, Philadelphia 3, Pa., U. S. A.

Just Published

New Classified Price Lists

Available separates from the TRANSACTIONS OF THE AMERICAN ENTOMOLOGICAL SOCIETY and ENTOMOLOGICAL NEWS, and all titles of the Society's MEMOIRS have been catalogued by author in twelve special price lists in the following categories:

Coleoptera	**Neuroptera and Smaller Orders**
Diptera	**Odonata**
Hemiptera	**Orthoptera-Dermaptera**
Hymenoptera	**Arachnida and Other Classes**
Lepidoptera	**Bibliography-Biography**
Memoirs	**General**

Lists will be mailed free upon request. Please state specifically which list or lists you require.

The American Entomological Society
1900 RACE STREET
PHILADELPHIA 3, PENNSYLVANIA

Just Published

MEMOIRS OF THE AMERICAN ENTOMOLOGICAL SOCIETY

Number 17

A TAXONOMIC STUDY OF THE MILLIPED FAMILY SPIROBOLIDAE (DIPLOPODA: SPIROBOLIDA)

By William T. Keeton

147 pages of text, 37 tables, 2 maps, 18 plates, table of contents and index

Spirobolid millipeds are probably the most widely known Diplopoda in the United States, being used in many college courses; yet the family has been little studied. This monograph brings together existing knowledge of the group for the first time, and adds much new information gained from critical study of series. The taxonomic history of the family is outlined. External morphology is briefly treated, with emphasis on characters utilized in classification. A summary of current knowledge of life histories is included. The family is redefined, and each genus and species is treated in detail. Particular attention is given to variation and distribution, both of which become more meaningful biologically as a result of synonymizing many species names. Possible phylogenetic relationships of the genera are discussed, and keys to all taxa are provided, with most diagnostic characters illustrated in 18 plates or summarized in 37 tables.

Price $5.50

THE AMERICAN ENTOMOLOGICAL SOCIETY

1900 Race Street, Philadelphia 3, Penna., U.S.A.